

# **EXPERIMENTAL TESTS FOR POLARIZED TIME INERTIA**

**David James Milton Hansen**

March 28, 2026

## **Abstract**

This companion document presents seven detailed experimental proposals for testing Polarized Time Inertia (PTI). Each test targets a specific prediction, providing theoretical background, experimental design, expected results under PTI vs. standard physics, required precision, and feasibility assessment.

## Test 1: Non-Linear Gravitational Superposition

### Theoretical Prediction

PTI predicts that gravitational fields superpose non-linearly due to recursive comparison structure.

$$g_{total} = \sum g_i + \epsilon * \sum_{i \neq j} g_i * g_j$$

### Experimental Setup

High-precision torsion balance or atom interferometer between two precisely known masses at separations 1 cm to 1 m. Measure gravitational acceleration at midpoint.

### Expected Results

**Standard physics:**  $g_{total} = g_1 + g_2$  exactly (linear superposition in weak-field GR).

**PTI prediction:**  $g_{total} = g_1 + g_2 + \epsilon(g_1 * g_2)$ , with  $\epsilon \sim 10^{-15}$ .

### Precision Required

Current sensitivity  $\sim 10^{-15}$  m/s<sup>2</sup>. Requires next-generation instruments.

### Feasibility

High. Apparatus exists. Main challenge: seismic noise and systematic control.

## Test 2: Higgs Field and Spatial Curvature Correlation

### Theoretical Prediction

If Higgs field = space, then Higgs boson properties should correlate with local spatial curvature.

$$\sigma_H(t) \text{ proportional to } (1 + \alpha * \Phi_{\text{tidal}}(t))$$

### Experimental Setup

Statistical analysis of LHC Run 2/3 Higgs events correlated with tidal gravitational potential variations from Moon/Sun ( $\Delta g \sim 10^{-7} \text{ m/s}^2$ ).

### Expected Results

**Standard physics:** No correlation between Higgs properties and local gravity.

**PTI prediction:** Small but statistically significant modulation of Higgs production rates with tidal cycles.

### Precision Required

Requires  $>10^5$  Higgs events. Available from LHC Run 2+3.

### Feasibility

Moderate. Data exists. Disentangling from accelerator performance systematics is challenging.

## Test 3: Entanglement in Varying Gravitational Fields

### Theoretical Prediction

PTI predicts entanglement correlations are subtly affected by gravitational field differences between detectors.

$$S_{CHSH} = 2\sqrt{2} * (1 + \beta * \Delta_{\Phi}/c^2)$$

### Experimental Setup

Bell test with polarization-entangled photons. One detector at ground, one at 10 km altitude (satellite or mountain). Compare CHSH parameter S for different configurations.

### Expected Results

**Standard physics:**  $S = 2\sqrt{2} = 2.828$  independent of gravitational potential.

**PTI prediction:** S deviation of order  $10^{-12}$  correlated with gravitational potential difference.

### Precision Required

Requires  $\sim 10^{24}$  entangled pairs for  $10^{-12}$  sensitivity.

### Feasibility

Moderate-high. Satellite entanglement experiments exist (Micius). Statistical precision is the challenge.

## Test 4: Dark Energy Equation of State Evolution

### Theoretical Prediction

PTI predicts  $w(z)$  is slightly evolving rather than exactly -1.

$$w(z) = -1 + \delta_0 * (\rho_m(z)/\rho_c)$$

### Experimental Setup

Constrain  $w(z)$  from Type Ia supernovae, BAO, and CMB power spectra. Compare PTI model vs. LCDM.

### Expected Results

**Standard physics:**  $w = -1$  exactly (cosmological constant).

**PTI prediction:**  $w(z)$  deviates from -1, with functional form predicted by PTI.

### Precision Required

Current surveys (DESI) constrain  $w$  to  $\sim 5\%$ . PTI needs to predict  $\delta_0$ .

### Feasibility

High. Data exists and is being analyzed. PTI must produce quantitative  $w(z)$  prediction.

## Test 5: Planck-Scale Lorentz Invariance Violation

### Theoretical Prediction

The discrete comparison structure implies energy-dependent light speed.

$$v(E) = c * (1 +/- (E/E_{Planck})^n), n = 1 \text{ or } 2$$

### Experimental Setup

Analyze GRB photon arrival times from Fermi telescope. Compare high-energy vs. low-energy photon arrival over cosmological distances.

### Expected Results

**Standard physics:** No energy-dependent arrival time differences.

**PTI prediction:** Systematic energy-arrival correlation. For  $n=1$ , current limits constrain  $E_{LIV} > 10^{19}$  GeV.

### Precision Required

Fermi data provides sensitivity to  $E_{LIV} \sim 10^{19}$  GeV for  $n=1$ .

### Feasibility

High. Data exists from Fermi, H.E.S.S., MAGIC, VERITAS.

## Test 6: Gravitational Wave Polarization Modes

### Theoretical Prediction

PTI's space-flow model may produce additional GW polarization modes beyond GR's two tensor modes.

$$h_{ij} = h_{+} * e_{+}^{ij} + h_{x} * e_{x}^{ij} + h_{scalar} * \delta^{ij}$$

### Experimental Setup

Multi-detector GW analysis (LIGO, Virgo, KAGRA) to decompose polarization content of detected events.

### Expected Results

**Standard physics:** Two tensor modes only (+ and x).

**PTI prediction:** Additional scalar breathing mode from space-flow oscillation.

### Precision Required

Requires 3+ detectors with different orientations.

### Feasibility

High. LIGO/Virgo/KAGRA network already operational.

## Test 7: Spatial Quantization in Tabletop Experiments

### Theoretical Prediction

Space is quantized at the Planck scale, detectable via accumulated effects.

$$\Delta x_{\min} = l_P = \sqrt{\hbar G/c^3} \sim 1.616 * 10^{-35} \text{ m}$$

### Experimental Setup

Tabletop optomechanical experiments measuring position uncertainty of a nanogram-scale mass at ultra-low temperatures.

### Expected Results

**Standard physics:** Position can be measured to arbitrary precision (limited only by Heisenberg).

**PTI prediction:** Position measurements show Planck-scale granularity, detectable as excess noise.

### Precision Required

Current tabletop sensitivity  $\sim 10^{-21}$  m. Needs improvement to  $10^{-35}$  m.

### Feasibility

Low-moderate. Far from Planck scale, but theoretical proposals exist for amplified signatures.

## Summary

Test	PTI Prediction	Standard	Feasibility
1. Grav. Nonlinearity	Non-linear	Linear	High
2. Higgs-Curvature	Correlated	No corr.	Moderate
3. Entangle. + Gravity	S deviation	S constant	Mod-High
4. Dark Energy $w(z)$	Evolving	$w=-1$	High
5. Lorentz Violation	E-dep. speed	Constant $c$	High
6. GW Polarization	Extra modes	2 modes	High
7. Spatial Quanta	Planck noise	No noise	Low-Mod

Each test targets specific distinguishing predictions. Several are feasible with current technology.

## References

- [1] A. Einstein, "Die Grundlage der allgemeinen Relativitätstheorie," *Annalen der Physik*, vol. 354, no. 7, pp. 769-822, 1916.
- [2] P. A. M. Dirac, "The Quantum Theory of the Electron," *Proceedings of the Royal Society A*, vol. 117, no. 778, pp. 610-624, 1928.
- [3] P. W. Higgs, "Broken Symmetries and the Masses of Gauge Bosons," *Physical Review Letters*, vol. 13, no. 16, pp. 508-509, 1964.
- [4] S. W. Hawking, "Particle Creation by Black Holes," *Communications in Mathematical Physics*, vol. 43, no. 3, pp. 199-220, 1975.
- [5] H. Everett III, "Relative State Formulation of Quantum Mechanics," *Reviews of Modern Physics*, vol. 29, no. 3, pp. 454-462, 1957.
- [6] J. S. Bell, "On the Einstein Podolsky Rosen Paradox," *Physics Physique Fizika*, vol. 1, no. 3, pp. 195-200, 1964.
- [7] R. P. Feynman, "Space-Time Approach to Non-Relativistic Quantum Mechanics," *Reviews of Modern Physics*, vol. 20, no. 2, pp. 367-387, 1948.
- [8] M. Planck, "Über das Gesetz der Energieverteilung im Normalspectrum," *Annalen der Physik*, vol. 309, no. 3, pp. 553-563, 1901.
- [9] K. Schwarzschild, "Über das Gravitationsfeld eines Massenpunktes," *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, pp. 189-196, 1916.
- [10] A. G. Riess et al., "Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant," *The Astronomical Journal*, vol. 116, no. 3, pp. 1009-1038, 1998.
- [11] S. Perlmutter et al., "Measurements of Omega and Lambda from 42 High-Redshift Supernovae," *The Astrophysical Journal*, vol. 517, no. 2, pp. 565-586, 1999.
- [12] V. C. Rubin, W. K. Ford Jr., "Rotation of the Andromeda Nebula from a Spectroscopic Survey of Emission Regions," *The Astrophysical Journal*, vol. 159, p. 379, 1970.
- [13] D. J. Gross, F. Wilczek, "Ultraviolet Behavior of Non-Abelian Gauge Theories," *Physical Review Letters*, vol. 30, no. 26, pp. 1343-1346, 1973.
- [14] ATLAS Collaboration, "Observation of a new particle in the search for the Standard Model Higgs boson," *Physics Letters B*, vol. 716, no. 1, pp. 1-29, 2012.
- [15] Planck Collaboration, "Planck 2018 results. VI. Cosmological parameters," *Astronomy & Astrophysics*, vol. 641, A6, 2020.
- [16] J. A. Wheeler, "Information, Physics, Quantum: The Search for Links," in *Proc. 3rd Int. Symp. Found. Quantum Mech.*, pp. 354-368, 1989.
- [17] L. Susskind, "The World as a Hologram," *Journal of Mathematical Physics*, vol. 36, no. 11, pp. 6377-6396, 1995.
- [18] R. Penrose, "On Gravity's Role in Quantum State Reduction," *General Relativity and Gravitation*, vol. 28, no. 5, pp. 581-600, 1996.
- [19] C. Rovelli, "Loop Quantum Gravity," *Living Reviews in Relativity*, vol. 11, no. 1, 2008.
- [20] J. Maldacena, "The Large N Limit of Superconformal Field Theories and Supergravity," *International Journal of Theoretical Physics*, vol. 38, no. 4, pp. 1113-1133, 1999.

- [21] A. Aspect, P. Grangier, G. Roger, "Experimental Tests of Bell's Inequalities Using Time-Varying Analyzers," *Physical Review Letters*, vol. 49, no. 25, pp. 1804-1807, 1982.
- [22] B. P. Abbott et al. (LIGO), "Observation of Gravitational Waves from a Binary Black Hole Merger," *Physical Review Letters*, vol. 116, no. 6, 061102, 2016.
- [23] D. J. M. Hansen, "Polarized Time Inertia: Foundational Notes," Unpublished research notes, August 2024 - March 2026.